



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2014

The sodium chloride cotransporter SLC12A3: new roles in sodium, potassium, and blood pressure regulation

Moes, Arthur D ; van der Lubbe, Nils ; Zietse, Robert ; Loffing, Johannes ; Hoorn, Ewout J

Abstract: SLC12A3 encodes the thiazide-sensitive sodium chloride cotransporter (NCC), which is primarily expressed in the kidney, but also in intestine and bone. In the kidney, NCC is located in the apical plasma membrane of epithelial cells in the distal convoluted tubule. Although NCC reabsorbs only 5 to 10 % of filtered sodium, it is important for the fine-tuning of renal sodium excretion in response to various hormonal and non-hormonal stimuli. Several new roles for NCC in the regulation of sodium, potassium, and blood pressure have been unraveled recently. For example, the recent discoveries that NCC is activated by angiotensin II but inhibited by dietary potassium shed light on how the kidney handles sodium during hypovolemia (high angiotensin II) and hyperkalemia. The additive effect of angiotensin II and aldosterone maximizes sodium reabsorption during hypovolemia, whereas the inhibitory effect of potassium on NCC increases delivery of sodium to the potassium-secreting portion of the nephron. In addition, great steps have been made in unraveling the molecular machinery that controls NCC. This complex network consists of kinases and ubiquitinases, including WNKs, SGK1, SPAK, Nedd4-2, Cullin-3, and Kelch-like 3. The pathophysiological significance of this network is illustrated by the fact that modification of each individual protein in the network changes NCC activity and results in salt-dependent hypotension or hypertension. This review aims to summarize these new insights in an integrated manner while identifying unanswered questions.

DOI: <https://doi.org/10.1007/s00424-013-1407-9>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-88126>

Journal Article

Accepted Version

Originally published at:

Moes, Arthur D; van der Lubbe, Nils; Zietse, Robert; Loffing, Johannes; Hoorn, Ewout J (2014). The sodium chloride cotransporter SLC12A3: new roles in sodium, potassium, and blood pressure regulation. *Pflügers Archiv : European Journal of Physiology*, 466(1):107-118.

DOI: <https://doi.org/10.1007/s00424-013-1407-9>

19 November 2013 – Revised version (changes in bold red font)

**The sodium chloride cotransporter SLC12A3:
new roles in sodium, potassium, and blood pressure regulation**

Arthur D. Moes¹, Nils van der Lubbe¹, Robert Zietse¹, Johannes Loffing^{2,3}, Ewout J. Hoorn¹

¹Department of Internal Medicine, Erasmus Medical Center, Rotterdam, The Netherlands

²Institute of Anatomy, University of Zürich, Zürich, Switzerland, ³Zurich Centre for Integrative
Human Physiology

Manuscript type : This article is published as part of the Special Issue on sodium-dependent
transporters in health and disease

Number of figures : 4

Number of tables : 2

Word count : 3884

Corresponding author:

Ewout J Hoorn, MD, PhD E-mail : e.j.hoorn@erasmusmc.nl

PO Box 2040 – Room H-438 Phone : +31-10-7032860

3000 CA Rotterdam, The Netherlands Fax : +31-10-7033008

Summary (word count: 228)

SLC12A3 encodes the thiazide-sensitive sodium chloride cotransporter (NCC), which is primarily expressed in the kidney, but also in intestine and bone. In the kidney, NCC is located in the apical plasma membrane of epithelial cells in the distal convoluted tubule. Although NCC reabsorbs only 5 to 10% of filtered sodium, it is important for the fine-tuning of renal sodium excretion in response to various hormonal and non-hormonal stimuli. Several new roles for NCC in the regulation of sodium, potassium, and blood pressure have been unraveled recently. For example, the recent discoveries that NCC is activated by angiotensin II but inhibited by dietary potassium sheds light on how the kidney handles sodium during hypovolemia (high angiotensin II) and hyperkalemia. The additive effect of angiotensin II and aldosterone maximizes sodium reabsorption during hypovolemia, whereas the inhibitory effect of potassium on NCC increases delivery of sodium to the potassium-secreting portion of the nephron. In addition, great steps have been made in unraveling the molecular machinery that controls NCC. This complex network consists of kinases and ubiquitinases, including WNKs, SGK1, SPAK, Nedd4-2, Cullin-3 and Kelch-like 3. The pathophysiological significance of this network is illustrated by the fact that modification of each individual protein in the network changes NCC activity and results in salt-dependent hypotension or hypertension. This review aims to summarize these new insights in an integrated manner while identifying unanswered questions.

Keywords: Aldosterone; Angiotensin II; Hypertension; Tacrolimus; Thiazide; WNK kinase.

Typical hallmarks of NCC

The presence of a sodium chloride cotransporter was first suggested in urinary bladder of the winter flounder [95, 96]. Subsequent studies in this tissue demonstrated that this cotransporter could be inhibited by thiazide diuretics [111] and the cDNA encoding this transporter was isolated [31].

Studies using micropuncture and isolated perfused tubules had identified the same pharmacological and kinetic transport characteristics in the early distal convoluted tubule (DCT) of rat kidney [23, 60]. Indeed, Gamba and colleagues succeeded in isolating cDNA of the sodium chloride cotransporter from rat kidney [30]. NCC is encoded by the *SLC12A3* gene (55-kb, 26 exons) and belongs to the SLC12 family of electroneutral cation chloride cotransporters [29].

Besides the kidney, NCC was shown to be expressed in intestine and bone where it is likely involved in sodium and calcium absorption [4, 20]. It has been suggested that NCC is expressed in various other tissues but this has not been confirmed [29]. In the kidney, NCC is located in the early part of the DCT (also called DCT1) (**Figure 1**), **but gradually decreases along the later** part of the DCT (DCT2), where it co-localizes with the epithelial sodium channel (ENaC) [3].

The sodium chloride cotransporter (NCC) mediates the reabsorption of sodium across the apical membrane of the DCT (**Figure 1**) [23]. Chronic activation or inhibition of NCC is usually accompanied by morphological changes in the DCT resulting in hypertrophy or atrophy [21, 61, 66, 126]. The gradient required for sodium chloride transport through NCC is generated and maintained by the basolateral sodium-potassium ATPase pump. The potassium that enters the cell via this pump is **recycled by basolateral potassium transporters. In addition, potassium may also be** secreted apically by the renal outer medullary potassium channel (ROMK) and a potassium chloride cotransporter [133]. In addition, NCC modulates transcellular magnesium and calcium reabsorption in the DCT through interaction with the transporters TRPM6 and

TRVP5, respectively (**Figure 1**) [16, 85]. Only 5 to 10% of the filtered load of sodium is reabsorbed in the DCT and this is primarily mediated by NCC [23]. Despite this modest contribution to overall sodium reabsorption, the NCC in the DCT together with ENaC in the connecting tubule (CNT) and the collecting duct (CD) fine-tunes the final concentration of sodium chloride in the urine. **This is possible** because it is not affected by the tubuloglomerular feedback [29]. Due to this **property**, NCC plays a pivotal role in extracellular fluid volume and blood pressure control [29].

Structure–function relationship

In humans, NCC is a membrane glycoprotein of 1,021-amino acid residues which resembles the general topology of the sodium-potassium-chloride cotransporters 1 and 2 (NKCC 1 and 2) [70]. NCC is able to form dimers, and it is likely that it functions as a dimer [18, 29]. It consists of 12 putative transmembrane (TM) spanning regions with a central hydrophobic domain (**Figure 2**). Between TM7 and TM8 there is a large extracellular hydrophilic loop that contains two glycosylation sites (N404 and N424) [50]. These glycosylation sites are essential for functioning and surface expression of NCC [40]. When glycosylation is eliminated, thiazide diuretics have much greater access to their binding sites, which suggests that glycosylation blocks the affinity for thiazides [50]. A study that interchanged the transmembrane regions between rat and flounder NCC revealed that affinity-modifying residues for chloride are located within the TM1-7 region [78]. **Especially**, a highly conserved glycine within TM4 plays a crucial role for the affinity of chloride [78]. The TM8-12 region, **distal to the extracellular loop**, is sensitive to thiazides. Site-directed mutagenesis was used to show that a single residue in TM11 defines the different affinity for thiazides between mammalian and flounder NCC [12]. Both **transmembrane regions** appear to have affinity for extracellular sodium. The central domain is flanked by a short amino-terminal domain (N-terminus) and a long carboxy-terminal domain (C-terminus) which are both located intracellularly (**Figure 2**) [29]. The N-terminus of NCC contains several conserved phosphorylation sites including threonine 46, 55 and 60 and serine 73 and 91 in humans [19]. In rats **and mice** these phosphorylation sites correspond to threonine 44, 53, 58 and serine 71, 89, **and 124**. Phosphorylation of NCC appears to determine its activity, especially at threonine 60 in humans or 58 in rat. Indeed, a mutation in threonine 60 is a common cause of Gitelman syndrome [65], the disorder resulting from NCC inactivation (see further). The

development of a knock-in mouse model with this mutation exhibits the Gitelman phenotype and did so because mutant NCC was restricted to the cytosol [138]. Because phosphorylated NCC has thus far only been found in the apical plasma membrane of the DCT, anchoring in the plasma membrane seems necessary for phosphorylation to occur [91]. This suggests that, apart from phosphorylation, the trafficking of NCC from subapical vesicles to the plasma membrane is also important [42].

Physiological functions

NCC is highly regulated by hormones, including aldosterone, angiotensin II, glucocorticoids, estrogen, insulin, norepinephrine, and vasopressin (**Table 1**) [13, 53, 82, 83, 91, 108, 118, 124, 125]. The fact that NCC is regulated by so many different hormones suggests that sodium reabsorption through NCC is an important homeostatic control mechanism. Aldosterone was the first hormone recognized to be capable of activating NCC [53] and the DCT is therefore part of what has been called the “aldosterone-sensitive distal nephron” (ASDN) [77], which was defined to comprise the DCT2, the CNT and the CD [67]. Experimentally, aldosterone upregulates NCC both when it is directly infused or when it is secreted in response to a low sodium diet [14, 53]. The acute effect of aldosterone only involves phosphorylation of NCC [56], whereas the chronic effect also increases the total protein abundance of NCC [53], which likely occurs independent from changes to NCC mRNA levels [1, 69, 119]. The regulation of NCC by aldosterone seems logical, because at least the endportion of the DCT expresses the enzyme 11-beta-hydroxysteroid dehydrogenase II which rapidly inactivates glucocorticoids and hence provides mineralocorticoid-sensitivity to the epithelial cells [6]. The discovery that angiotensin II is also capable of activating NCC was more surprising, because its actions were believed to be confined to the proximal tubule [75]. By adrenalectomizing rats and selectively re-infusing aldosterone or angiotensin II, we were able to dissect the stimulatory effects of angiotensin II and aldosterone on NCC [118]. In a subsequent study we showed that aldosterone did not require angiotensin II for activation of NCC, although the presence of both stimuli led to an additive response [119]. This additive response may be useful to maximize sodium reabsorption during hypovolemia, because this is characterized by elevated plasma levels of both angiotensin II and aldosterone [122]. Increased sodium reabsorption in the DCT by NCC will also decrease the delivery of

sodium to the CNT and CD [42]. Because in these segments sodium reabsorption is electrochemically coupled to potassium secretion, decreased delivery of sodium will help conserve potassium. In line with this, recent studies have shown that angiotensin II directly inhibits ROMK, further contributing to potassium conservation during hypovolemia [49, 129, 140]. The effects of angiotensin II on NCC and ROMK therefore help to understand the aldosterone paradox, the question how aldosterone increases sodium reabsorption during hypovolemia, but potassium secretion during hyperkalemia [3, 122]. Further insight into the aldosterone paradox comes from the effects of potassium on NCC. Recent studies have shown that dietary potassium inhibits NCC [110, 121]. Gastric gavage of potassium in mice increased both urinary potassium and sodium excretion [110]. This response occurred within minutes and was independent from aldosterone, because it could also be induced in aldosterone-deficient animals. In the kidney, this was accompanied by dephosphorylation of NCC (**Figure 4A**). The increased delivery of sodium to the more distal nephron parts facilitates sodium–potassium exchange and therefore kaliuresis. This “potassium-induced natriuresis” appears to constitute an important physiological response. Namely, potassium-induced natriuresis could still be evoked when a high potassium diet was combined with a low sodium diet [121]. This was accompanied by a reduced abundance of total NCC; **phosphorylated NCC was also reduced, although this did not reach statistical significance** (**Figure 4B**). Taken together, these **and other** recent results seem to suggest that when the organism **is** faced with the choice between conserving sodium or secreting potassium, it chooses the latter [27]. The focus of future research will be to identify the signal by which dietary potassium induces NCC downregulation. Similar to high dietary potassium, high dietary sodium also suppresses NCC [14]. This response, however, is not as rapid as the one for high potassium diet [110] and involves a decrease in the plasma

membrane abundance of sodium transporters all along the nephron [28, 137]. Furthermore, the effect of high dietary sodium on NCC appears to be mediated through aldosterone [14], **although the effect of high dietary sodium on NCC has not been studied in the absence of aldosterone.** Similarly, chronic metabolic acidosis also increases aldosterone and therefore NCC [26]. Why other hormones such as insulin [13, 57, 108, 109], vasopressin [83, 91, 104], estrogen [125], and norepinephrine [82] regulate NCC is less clearly defined (**Table 1**), but warrants further study given the role of these hormones in normal physiology and human diseases such as diabetes mellitus, obesity, and hypertension.

The NCC signaling cascade

Kinases

The intracellular signaling cascade that controls NCC activity has largely been unraveled in recent years (**Figure 3**). This NCC signaling cascade consists of a multikinase network which includes the kinases WNK, SPAK, OSR1, and SGK1 [130]. More recently, proteins involved in ubiquitylation including Nedd4-2, Kelch-like 3, and Cullin 3 were also found to regulate NCC [2, 7, 68, 99]. Many of these regulatory proteins were identified because mutations in their genes result in familial hyperkalemic hypertension (FHHt, also called pseudohypoaldosteronism type II or Gordon syndrome, see also “Relation to diseases” below). WNKs **appear to** modulate both the “trafficking” and phosphorylation of NCC [41], **although most of our knowledge has been derived from studies in oocytes.** The regulation of NCC trafficking by WNKs involves a sequential inhibitory cascade, in which KS-WNK1 inhibits WNK1, WNK1 inhibits WNK4 [136], and WNK4 inhibits NCC [134]. The inhibition of NCC by WNK4 is not caused by endocytosis [35], but rather by promoting lysosomal degradation [112]. This inhibition is mediated by the ERK1/2 signaling pathway and the lysosomal targeting receptor sortilin [141],

142]. Interestingly, angiotensin II converts WNK4 from an inhibitor to an activator of NCC [101]. In contrast to WNK4, WNK3 stimulates NCC [97], but its actions are less well-defined. WNK3 and WNK4 not only have divergent effects on NCC, they also antagonize each other. Indeed, it appears to be the ratio between WNK3 and WNK4 that determines the net effect on NCC [135]. The phosphorylation of NCC is mediated by SPAK [79]; several WNKs interact with SPAK and therefore indirectly control the phosphorylation-step of NCC. Interactions between SPAK and WNK1 [79], WNK3 [34], and WNK4 [102] have been reported. However, the brain but not the kidney isoform of WNK3 can activate NCC and does so through a SPAK-independent mechanism [34]. Two isoforms of SPAK have been identified, including full-length SPAK (FL-SPAK) and kidney-specific SPAK (KS-SPAK or SPAK2), of which the latter isoform has low expression levels in the DCT [73]. KS-SPAK, which lacks the kinase domain, inhibits FL-SPAK and OSR1, which are both known to phosphorylate NKCC2. This may explain why in mice the knockout of SPAK results in decreased NCC phosphorylation (absence of full-length SPAK), but increased NKCC2 phosphorylation (no inhibition of FL-SPAK or OSR1 by KS-SPAK) [36, 73].

SPAK deficiency, however, does not completely inhibit NCC phosphorylation [73, 139], suggesting the involvement of other kinases or phosphatases. Both SPAK isoforms are also involved in the stimulatory effect of vasopressin on NCC and NKCC2 [83, 91]. Namely, vasopressin stimulates FL-SPAK in the DCT to phosphorylate NCC, whereas it attenuates KS-SPAK to allow FL-SPAK and OSR1 to phosphorylate NKCC2 [104]. Although SGK1 was first recognized as an activator of ENaC [127], later reports also showed effects on NCC [115]. SGK1 and NCC do not seem to interact directly, but rather through WNK4 and Nedd4-2 [2, 98]. SGK1 phosphorylates WNK4 and this phosphorylation step reduces the inhibition of WNK4 on NCC

[98, 100]. Because SGK1 is sensitive to aldosterone, this pathway appears to be involved in the activation of NCC by aldosterone [100]. The opposite is also true, because SGK1 knockout mice failed to increase NCC activity during a low sodium diet [115].

Ubiquitin ligases

Recent data indicate that Nedd4-2 is yet another player in the pathway by which aldosterone activates NCC (**Figure 3**) [2, 99]. Nedd4-2 was shown to stimulate ubiquitylation of NCC and decreased its activity and surface expression *in vitro* and *in vivo*, while SGK1 prevented these effects [2]. The pathophysiological significance of the regulation of NCC by Nedd4-2 was shown by the generation of inducible nephron-specific Nedd4-2 knockout mice [99]. These mice exhibited salt-dependent hypertension that was characterized by upregulation of total and phosphorylated NCC and sensitivity to thiazides. The deletion of Nedd4-2 also affected ENaC and ROMK, which were down- and upregulated, respectively. This may explain the additional characteristics of these mice, namely that they had a normal Na^+/K^+ balance and were not hyperkalemic. Similar to Nedd4-2, Kelch-like 3 and Cullin 3 are also involved in ubiquitylation, because they are components of an E3 ubiquitin ligase complex. Although mutations in Kelch-like 3 and Cullin-3 cause hyperkalemic hypertension that is reversible with thiazides, these proteins probably do not interact directly with NCC. Instead, Kelch-like 3 binds and ubiquitinates WNK4 and the subsequent degradation of WNK4 would be expected to increase NCC activity (**Figure 3**) [89, 106, 128, 132].

Other signaling molecules

Although the network reviewed above is largely interconnected (**Figure 3**), additional NCC regulatory pathways exist. One example is the regulation of NCC by phorbol esters [54]. This

effect is mediated through Ras guanyl-releasing protein 1 and ERK1/2, which stimulates ubiquitylation and endocytosis of NCC [55]. Furthermore, NCC phosphorylation is not only controlled by kinases but also by phosphatases including phosphatase 4 [33]. Finally, the process that controls the phosphorylation of NCC may also cause less ubiquitylation, thereby increasing the number of cotransporters available for phosphorylation [51].

Relation to human disease

Gitelman syndrome

The clearest demonstration of the functional relevance of NCC comes from human monogenetic diseases that affect NCC function (**Table 2**). Inactivating mutations in *SLC12A3* cause the so-called Gitelman syndrome [107]. Gitelman syndrome is an autosomal recessive disorder that is characterized by hypokalemia, hypomagnesemia, metabolic alkalosis, hypocalciuria, and low to normal arterial blood pressure. Missense mutations account for approximately 59% of the mutations in Gitelman syndrome and compound heterozygosity is common [113, 123]. Gender and the type of mutation contribute to phenotypic variability, with males and patients with homozygous deep intronic mutations exhibiting a **more severe** phenotype [64, 65]. Novel mutations are still being identified in Gitelman syndrome and these genetic defects lead to impaired production, processing, insertion or regulation of the NCC protein (type I, II, III or IV mutations) [32, 113, 123]. Although Gitelman syndrome is usually a relatively benign disorder that often remains subclinical for many years, a recent report suggests that the electrolyte disorders associated with Gitelman syndrome may result in chronic kidney disease and glucose intolerance [113]. On the other hand, heterozygous mutations in NCC may prevent hypertension and cardiovascular diseases [52] and improve bone-density [17] likely because they induce mild sodium excretion and a positive calcium balance. NCC knockout mice recapitulate the phenotype of Gitelman syndrome, although some features become manifest only when the animals are challenged [73, 80, 105, 139]. In line with the current model of NCC regulation (**Figure 3**), genetically modified mice with overexpression of WNK4, deficiency in SPAK or its kinase domain, also exhibit features of Gitelman syndrome [61, 88, 94, 139], although such mutations have not been identified in humans. Gitelman syndrome should be differentiated from the more

severe Bartter syndrome, which results from mutations affecting NKCC2, and is characterized by earlier onset of symptoms and hypercalciuria instead of hypocalciuria [47]. An intermediary phenotype is caused by mutations in *CLCNKB* which encodes a basolateral chloride channel that is expressed in both the thick ascending limb and the DCT (**Table 2**) [58]. **Disturbed basolateral chloride efflux will indirectly impair apical NaCl transport through NKCC2 and NCC.**

Familial hyperkalemic hypertension

Familial hyperkalemic hypertension (FHHT) is the “mirror image” of Gitelman syndrome because, in addition to hyperkalemia and hypertension, it is characterized by hypercalciuria and metabolic acidosis [37]. Surprisingly, no activating mutations in *SLC12A3* have been reported. Overexpression of NCC in transgenic mice also failed to induce hyperkalemic hypertension, but this may have been due to the fact that phosphorylated NCC was not increased [74]. Similarly, increasing NCC activity by inactivating KS-WNK1 was also not sufficient to cause hyperkalemic hypertension, because it was associated with a compensatory decrease in ENaC [38]. Instead, familial hyperkalemic hypertension is caused by mutations in the genes encoding WNK1, WNK4, Kelch-like 3 or Cullin-3, which result in overactivity of NCC (**Table 2**) [7, 68, 131]. Because the WNKs regulate other transporters than NCC alone, these effects may also contribute to the FHHT-phenotype [37]. Intronic deletions in the WNK1 gene cause overexpression of WNK1 and therefore more inhibition of WNK4. The inhibition of WNK4 will relieve the inhibition of SPAK and will activate NCC (**Figure 3**). Similarly, missense mutations in the WNK4 gene give rise to a mutant protein that no longer inhibits SPAK resulting in NCC activation. Mutations in *KLHL3* can be dominant or recessive and homozygous or heterozygous,

1 while the identified mutations in *CUL3* were dominant, heterozygous and often *de novo* [7].
2
3
4
5
6 Mutations in *KLHL3* and *CUL3* seem to abrogate ubiquitylation of targets normally bound by
7
8
9 *KLHL3* including WNK4 [106]. Of the different mutations causing FHHt, *CUL3* mutations have
10
11 the most severe phenotype, including the youngest onset of hypertension and the highest serum
12
13 potassium [7]. Another illustration that NCC activation is the final common pathway of the
14
15 FHHt mutations is the fact that these disorders are all exquisitely sensitive to treatment with
16
17 thiazide-type diuretics [71].
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Drugs influencing NCC activity

Drugs inhibiting NCC

Several commonly used drugs inhibit or stimulate NCC activity (**Table 1**). The sensitivity of NCC for thiazide diuretics is well known and characterizes this cotransporter. Thiazides, however, do not exclusively inhibit NCC, but also the sodium-dependent bicarbonate-chloride cotransporter in the cortical collecting duct [63]. Thiazide diuretics were incidentally discovered while searching for better carbonic anhydrase inhibitors [86]. They were implemented clinically in 1958, long before it became apparent that their primary target is NCC [81]. Thiazide diuretics are the logical drug of choice for diseases with NCC overactivity such as the rare disorder FHHt. However, the clinical indication for thiazide diuretics is much broader. In fact, thiazide diuretics are still among the most commonly used drugs to treat hypertension worldwide. Although the natriuretic effect of these drugs undoubtedly contributes to their antihypertensive effect, this response also seems to be translated to an effect on vascular tone [9-11]. In addition, thiazides may directly cause vasodilation possibly through vascular potassium channel activation [92]. In addition to hypertension, thiazide diuretics can also be used for sodium-retaining disorders such as heart failure, liver cirrhosis, nephrotic syndrome, and chronic kidney disease [8]. Because thiazide diuretics act in the tubular lumen, their action requires successful delivery to the DCT. This is mediated by active secretion of thiazide diuretics in the proximal tubule through organic anion transporter and multidrug resistance-associated protein 4 [39, 114]. This explains why a reduction in glomerular filtration rate results in reduced efficacy of thiazide diuretics. Patients with Gitelman syndrome also demonstrate a dramatically impaired natriuretic response to thiazides, a feature that may be used diagnostically [15]. The side-effect profile of thiazide diuretics resembles Gitelman syndrome with the exception of hyponatremia, which is only seen with thiazides [22, 24, 48].

Drugs stimulating NCC

Drugs *stimulating* NCC activity have also been identified recently and they include the calcineurin inhibitors cyclosporine and tacrolimus (**Table 1**) [44, 76]. Calcineurin inhibitors are potent immunosuppressive drugs that are used clinically to prevent rejection after transplantation and sometimes in autoimmune disorders. Although cyclosporine and tacrolimus have different intracellular binding proteins, they both activate NCC, and this effect therefore appears to be a class effect [45]. Again, cyclosporine and tacrolimus do not seem to activate NCC directly, but influence the WNKs [76]. This possibility was already suggested by the side-effect profile of these drugs, which is similar to the phenotype of FHHt. Of interest, calcineurin is a protein phosphatase and protein phosphatases were recently shown to regulate NCC [33]. We were able to illustrate the clinical relevance of NCC activation by tacrolimus by linking it to hypertension [44]. That is, tacrolimus failed to induce hypertension in NCC knockout mice, whereas it caused more severe hypertension in NCC transgenic mice. Furthermore, a thiazide diuretic caused a larger urinary chloride excretion in patients on tacrolimus than in healthy volunteers or patients on sirolimus (a different immunosuppressant). This enhanced chloriuretic response to a thiazide was interpreted as an indication of increased transporter activity [15]. Also, the expression of total and phosphorylated NCC was increased in the kidney biopsies of patients treated with tacrolimus. Of interest, a model of cyclosporine nephrotoxicity showed the opposite effect on NCC, but this was attributed to inactivation of the renin-angiotensin system [62]. Furosemide, which blocks sodium transport in the thick ascending limb but not in the DCT, also increases NCC abundance. This effect is likely caused by the loop-diuretic induced enhanced sodium delivery to the DCT and the activation of the renin-angiotensin system [1]. The up-regulation of NCC likely compensates for the furosemide effect and hence may contribute to the phenomenon

of loop-diuretic resistance which frequently develops during chronic treatment. As such it provides the rationale for the combination of a loop-diuretic with a thiazide to overcome diuretic resistance [8]. Furthermore, recent studies in rats suggested that the cytostatic drug cisplatin decreases renal NCC abundance. However, this effect is likely attributed to a general toxic effect on DCT cells, because also the expression of other DCT-specific proteins such as the magnesium channel TRPM6 and the calcium- and magnesium-binding protein parvalbumin were decreased [116].

Perspectives

In this review we have highlighted several exciting new roles of NCC in sodium, potassium, and blood pressure regulation. These new insights have partly solved longstanding questions in physiology, but at the same time raise new questions. For example, although the model of NCC regulation is gaining more clarity (**Figure 3**), some of the interactions are still not well understood or even controversial. For example, WNK4 seems to be a negative regulator of NCC under some conditions, but may become a positive regulator in others [72]. The inhibition of NCC by dietary potassium may be mediated by WNK4 [87, 110, 120], but leaves the question open through which signal DCT cells “sense” dietary potassium. Pathophysiologically, the role of NCC in “essential” hypertension will likely remain a focus of future studies. The recent linkage of Cullin-3 and Kelch-like 3 to FHHt begs the question whether polymorphisms in these genes exist and whether they may contribute to human hypertension. The discovery that calcineurin inhibitors stimulate NCC to cause hypertension warrants a clinical study to evaluate whether thiazide diuretics are effective drugs to treat this side effect [44-46]. Especially for translational studies it will be important to have a measure of NCC activity *in vivo*. In addition to testing the response to a thiazide diuretic [15], the analysis of NCC in so-called urinary exosomes holds promise [43]. Urinary exosomes are vesicles derived from renal tubular epithelial cells that are thought to reflect the metabolic profile of these cells [93]. We recently showed that the abundance of phosphorylated NCC in urinary exosomes correlated with elevated aldosterone levels in animals and humans [117]. Although this review focused on NCC, sodium excretion by the kidney depends on many other sodium transporters, including NHE3, NKCC2, ENaC, and pendrin. Stimuli that activate NCC, sometimes also activate these other transporters, but an opposite, compensatory response may also occur [38, 84, 99]. In conclusion, the role of NCC in normal physiology and in the pathophysiology of hypertension is expanding and will

likely continue to do so in coming years. To fully grasp the potential of these insights for the treatment of human disease will likely require a more complete understanding of the molecular physiology of this fascinating cotransporter.

Grants

EJH is supported by the Dutch Kidney Foundation (KJPB 08.004) and the Netherlands

Organisation for Scientific Research (Veni). NvdL is supported by the Netherlands Foundation for Cardiovascular Excellence. JL is supported by the Swiss National Science Foundation, the Swiss National Centre in Competence in Research Kidney.CH, and the Zurich Centre for Integrative Human Physiology.

Table 1: Hormones, metabolic stimuli and drugs influencing NCC activity

	Stimulus	Effect	References
Hormones	Angiotensin II	Stimulatory	[118]
	Aldosterone	Stimulatory¶	[53]
	Glucocorticoids	Stimulatory	[124]
	Vasopressin	Stimulatory	[83, 91, 104]
	Insulin	Stimulatory	[13, 57, 108, 109]
	Estrogen	Stimulatory	[125]
	Norepinephrine	Stimulatory	[82]
Metabolic stimuli	Dietary potassium	Inhibitory	[110, 120]
	Dietary sodium*	Inhibitory	[14, 103]
	Dietary magnesium*	Stimulatory	[25]
	Acidosis*	Stimulatory	[26]
Drugs	Thiazide diuretics	Inhibitory	[16, 23]
	Furosemide*	Stimulatory	[1]
	Tacrolimus	Stimulatory	[44]
	Cyclosporine†	Stimulatory	[76]
	Cisplatin	Inhibitory	[116]

Footnote: ¶ When high dietary K increases aldosterone, NCC may be inhibited; * Some of these effects may be mediated through aldosterone; † In a model of cyclosporine nephrotoxicity,

NCC was downregulated, but this may be attributed to kidney failure and reduced renin-angiotensin activity [76].

Table 2: Characteristics of mutations causing Gitelman syndrome or familial hyperkalemic hypertension

Disease	Gene	OMIM gene	Human chromosome location	Encoding protein	Major phenotype	Selected references
Gitelman (OMIM 263800)	<i>SLC12A3</i>	600968	16q13	NCC	Hypokalemic hypotension	[107]
	<i>CLCNKB</i>	602023	1p36.13	CLCNKB	Variable	[58]
FHHt* (OMIM 179820)	<i>WNK1</i>	605232	12p13	WNK1	Hyperkalemic hypertension	[131]
	<i>WNK4</i>	601844	17q21	WNK4	Hyperkalemic hypertension	[131]
	<i>KLHL3</i>	605775	5q31	Kelch- like 3	Hyperkalemic hypertension	[7]
	<i>CUL-3</i>	603136	2q36	Cullin-3	Hyperkalemic hypertension	[7, 68]

Abbreviation: * FHHt, familial hyperkalemic hypertension; OMIM, online Mendelian inheritance in man.

Figure legends

Figure 1: Model of transcellular transport in the early distal convoluted tubule (DCT). A kidney tubule is shown schematically on the left indicating the locations of DCT type 1 and type 2 (DCT1, DCT2) and the connecting tubule (CNT). The sodium chloride cotransporter (NCC) is primarily expressed in DCT1. A model of transcellular transport in DCT1 is shown on the right, including the apical transporters NCC and transient receptor potential channels TRPV5 (a calcium channel) and TRPM6 (a magnesium channel). On the basolateral side the sodium potassium ATPase pump is shown as well as the chloride channel ClC-Kb and the sodium-calcium channel NCX1. This figure was adapted from [5, 59].

Figure 2: Putative structure of the sodium chloride cotransporter. The twelve transmembrane domains are shown including the hydrophilic loop with the two glycosylation sites. A detailed image of the N-terminus is provided on the left showing the binding sites of γ -adducin and the kinases SPAK/OSR1. This figure was reproduced and adapted from [19] with kind permission.

Figure 3: Current model of sodium chloride cotransporter regulation by kinases and ubiquitinases. The various interactions of the NCC regulatory pathway are shown as arrows (stimulatory) or as lines ending with perpendicular lines (inhibitory). Phosphorylation is indicated with the symbol “P”, whereas ubiquitylation is shown as “U”. SPAK/OSR1, WNK4, kidney-specific WNK1 (KS-WNK1) and long WNK1 (L-WNK1, also called WNK1) are kinases. The role of mutant WNK4 in familial hyperkalemic hypertension (FHHt) is also shown, which overrides the inhibitory effect of wild-type WNK4 on SPAK/OSR1. Nedd4-2 is a ubiquitinase, while Cullin-3 and Kelch-like 3 interact in a ubiquitylation complex that likely

ubiquitinates WNK4. **Although WNK3 has been shown to interact with WNK4 and SPAK [90, 135], its precise role in NCC regulation remains less clear and we therefore decided not to include it.** See text for further details.

Figure 4: Inhibition of the sodium chloride cotransporter by dietary potassium (**“high K”**). The results of our recent studies on the inhibitory effect of dietary potassium on NCC are shown [110, 121]. Panel A shows that dietary potassium acutely downregulates phosphorylated NCC but not total NCC. Conversely, panel B shows that a chronic high potassium diet primarily decreased total NCC, but phosphorylated NCC less so. **The high potassium diet was 2% and 5%, respectively, and the low sodium diet was < 0.001% [110, 121].**

References

1. Abdallah JG, Schrier RW, Edelstein C, Jennings SD, Wyse B, Ellison DH (2001) Loop diuretic infusion increases thiazide-sensitive Na(+)/Cl(-)-cotransporter abundance: role of aldosterone. *J Am Soc Nephrol* 12:1335-1341
2. Arroyo JP, Lagnaz D, Ronzaud C, Vazquez N, Ko BS, Moddes L, Ruffieux-Daidie D, Hausel P, Koesters R, Yang B, Stokes JB, Hoover RS, Gamba G, Staub O (2011) Nedd4-2 modulates renal Na⁺-Cl⁻ cotransporter via the aldosterone-SGK1-Nedd4-2 pathway. *J Am Soc Nephrol* 22:1707-1719
3. Arroyo JP, Ronzaud C, Lagnaz D, Staub O, Gamba G (2011) Aldosterone paradox: differential regulation of ion transport in distal nephron. *Physiology (Bethesda)* 26:115-123
4. Bazzini C, Vezzoli V, Sironi C, Dossena S, Ravasio A, De Biasi S, Garavaglia M, Rodighiero S, Meyer G, Fascio U, Furst J, Ritter M, Botta G, Paulmichl M (2005) Thiazide-sensitive NaCl-cotransporter in the intestine: possible role of hydrochlorothiazide in the intestinal Ca²⁺ uptake. *J Biol Chem* 280:19902-19910
5. Bindels RJ (2010) 2009 Homer W. Smith Award: Minerals in motion: from new ion transporters to new concepts. *J Am Soc Nephrol* 21:1263-1269
6. Bostanjoglo M, Reeves WB, Reilly RF, Velazquez H, Robertson N, Litwack G, Morsing P, Dorup J, Bachmann S, Ellison DH (1998) 11 β -hydroxysteroid dehydrogenase, mineralocorticoid receptor, and thiazide-sensitive Na-Cl cotransporter expression by distal tubules. *J Am Soc Nephrol* 9:1347-1358
7. Boyden LM, Choi M, Choate KA, Nelson-Williams CJ, Farhi A, Toka HR, Tikhonova IR, Bjornson R, Mane SM, Colussi G, Lebel M, Gordon RD, Semmekrot BA, Poujol A, Valimaki MJ, De Ferrari ME, Sanjad SA, Gutkin M, Karet FE, Tucci JR, Stockigt JR, Keppler-Noreuil KM, Porter CC, Anand SK, Whiteford ML, Davis ID, Dewar SB, Bettinelli A, Fadrowski JJ, Belsha CW, Hunley TE, Nelson RD, Trachtman H, Cole TR, Pinsk M, Bockenhauer D, Shenoy M, Vaidyanathan P, Foreman JW, Rasoulpour M, Thameem F, Al-Shahrouri HZ, Radhakrishnan J, Gharavi AG, Goilav B, Lifton RP (2012) Mutations in kelch-like 3 and cullin 3 cause hypertension and electrolyte abnormalities. *Nature* 482:98-102
8. Brater DC (1998) Diuretic therapy. *N Engl J Med* 339:387-395
9. Calo L, Davis PA, Semplicini A (2002) Reduced content of alpha subunit of Gq protein content in monocytes of Bartter and Gitelman syndromes: relationship with vascular hyporeactivity. *Kidney Int* 61:353-354
10. Calo LA, Davis PA (2010) Number and function of circulating endothelial progenitor cells and calcitonin gene-related peptide in hypertension: support from and opportunities in Bartter's and Gitelman's syndromes patients. *J Hypertens* 28:2169-2170; author reply 2171
11. Calo LA, Puato M, Schiavo S, Zanardo M, Tirrito C, Pagnin E, Balbi G, Davis PA, Palatini P, Pauletto P (2008) Absence of vascular remodelling in a high angiotensin-II state (Bartter's and Gitelman's syndromes): implications for angiotensin II signalling pathways. *Nephrol Dial Transplant* 23:2804-2809
12. Castaneda-Bueno M, Vazquez N, Bustos-Jaimes I, Hernandez D, Rodriguez-Lobato E, Pacheco-Alvarez D, Carino-Cortes R, Moreno E, Bobadilla NA, Gamba G (2010) A single residue in transmembrane domain 11 defines the different affinity for thiazides

- between the mammalian and flounder NaCl transporters. *Am J Physiol Renal Physiol* 299:F1111-1119
13. Chavez-Canales M, Arroyo JP, Ko B, Vazquez N, Bautista R, Castaneda-Bueno M, Bobadilla NA, Hoover RS, Gamba G (2013) Insulin increases the functional activity of the renal NaCl cotransporter. *J Hypertens* 31:303-311
 14. Chiga M, Rai T, Yang SS, Ohta A, Takizawa T, Sasaki S, Uchida S (2008) Dietary salt regulates the phosphorylation of OSR1/SPAK kinases and the sodium chloride cotransporter through aldosterone. *Kidney Int* 74:1403-1409
 15. Colussi G, Bettinelli A, Tedeschi S, De Ferrari ME, Syren ML, Borsa N, Mattiello C, Casari G, Bianchetti MG (2007) A thiazide test for the diagnosis of renal tubular hypokalemic disorders. *Clin J Am Soc Nephrol* 2:454-460
 16. Costanzo LS (1985) Localization of diuretic action in microperfused rat distal tubules: Ca and Na transport. *Am J Physiol* 248:F527-535
 17. Cruz DN (2001) The renal tubular Na-Cl co-transporter (NCCT): a potential genetic link between blood pressure and bone density? *Nephrol Dial Transplant* 16:691-694
 18. de Jong JC, Willems PH, Mooren FJ, van den Heuvel LP, Knoers NV, Bindels RJ (2003) The structural unit of the thiazide-sensitive NaCl cotransporter is a homodimer. *J Biol Chem* 278:24302-24307
 19. Dimke H (2011) Exploring the intricate regulatory network controlling the thiazide-sensitive NaCl cotransporter (NCC). *Pflugers Arch* 462:767-777
 20. Dvorak MM, De Joussineau C, Carter DH, Pisitkun T, Knepper MA, Gamba G, Kemp PJ, Riccardi D (2007) Thiazide diuretics directly induce osteoblast differentiation and mineralized nodule formation by interacting with a sodium chloride co-transporter in bone. *J Am Soc Nephrol* 18:2509-2516
 21. Ellison DH (2012) Adaptation in Gitelman syndrome: "we just want to pump you up". *Clin J Am Soc Nephrol* 7:379-382
 22. Ellison DH, Loffing J (2009) Thiazide effects and adverse effects: insights from molecular genetics. *Hypertension* 54:196-202
 23. Ellison DH, Velazquez H, Wright FS (1987) Thiazide-sensitive sodium chloride cotransport in early distal tubule. *Am J Physiol* 253:F546-554
 24. Ernst ME, Moser M (2009) Use of diuretics in patients with hypertension. *N Engl J Med* 361:2153-2164
 25. Fanestil DD, Hyde RH, Blakely P, Vaughn DA (1999) Dietary magnesium, not calcium, regulates renal thiazide receptor. *J Am Soc Nephrol* 10:458-463
 26. Farouqi S, Sheriff S, Amlal H (2006) Metabolic acidosis has dual effects on sodium handling by rat kidney. *Am J Physiol Renal Physiol* 291:F322-331
 27. Frindt G, Houde V, Palmer LG (2011) Conservation of Na⁺ vs. K⁺ by the rat cortical collecting duct. *Am J Physiol Renal Physiol* 301:F14-20
 28. Frindt G, Palmer LG (2009) Surface expression of sodium channels and transporters in rat kidney: effects of dietary sodium. *Am J Physiol Renal Physiol* 297:F1249-1255
 29. Gamba G (2009) The thiazide-sensitive Na⁺-Cl⁻ cotransporter: molecular biology, functional properties, and regulation by WNKs. *Am J Physiol Renal Physiol* 297:F838-848
 30. Gamba G, Miyanoshita A, Lombardi M, Lytton J, Lee WS, Hediger MA, Hebert SC (1994) Molecular cloning, primary structure, and characterization of two members of the

- mammalian electroneutral sodium-(potassium)-chloride cotransporter family expressed in kidney. *J Biol Chem* 269:17713-17722
31. Gamba G, Saltzberg SN, Lombardi M, Miyanoshita A, Lytton J, Hediger MA, Brenner BM, Hebert SC (1993) Primary structure and functional expression of a cDNA encoding the thiazide-sensitive, electroneutral sodium-chloride cotransporter. *Proc Natl Acad Sci U S A* 90:2749-2753
 32. Glaudemans B, Yntema HG, San-Cristobal P, Schoots J, Pfundt R, Kamsteeg EJ, Bindels RJ, Knoers NV, Hoenderop JG, Hoefsloot LH (2012) Novel NCC mutants and functional analysis in a new cohort of patients with Gitelman syndrome. *Eur J Hum Genet* 20:263-270
 33. Glover M, Mercier Zuber A, Figg N, O'Shaughnessy KM (2010) The activity of the thiazide-sensitive Na(+)-Cl(-) cotransporter is regulated by protein phosphatase PP4. *Can J Physiol Pharmacol* 88:986-995
 34. Glover M, Zuber AM, O'Shaughnessy KM (2009) Renal and brain isoforms of WNK3 have opposite effects on NCCT expression. *J Am Soc Nephrol* 20:1314-1322
 35. Golbang AP, Cope G, Hamad A, Murthy M, Liu CH, Cuthbert AW, O'Shaughnessy K M (2006) Regulation of the expression of the Na/Cl cotransporter by WNK4 and WNK1: evidence that accelerated dynamin-dependent endocytosis is not involved. *Am J Physiol Renal Physiol* 291:F1369-1376
 36. Grimm PR, Taneja TK, Liu J, Coleman R, Chen YY, Delpire E, Wade JB, Welling PA (2012) SPAK isoforms and OSR1 regulate sodium-chloride co-transporters in a nephron-specific manner. *J Biol Chem* 287:37673-37690
 37. Hadchouel J, Delaloy C, Faure S, Achard JM, Jeunemaitre X (2006) Familial hyperkalemic hypertension. *J Am Soc Nephrol* 17:208-217
 38. Hadchouel J, Soukaseum C, Busst C, Zhou XO, Baudrie V, Zurrer T, Cambillau M, Elghozi JL, Lifton RP, Loffing J, Jeunemaitre X (2010) Decreased ENaC expression compensates the increased NCC activity following inactivation of the kidney-specific isoform of WNK1 and prevents hypertension. *Proc Natl Acad Sci U S A* 107:18109-18114
 39. Hasegawa M, Kusuhara H, Adachi M, Schuetz JD, Takeuchi K, Sugiyama Y (2007) Multidrug resistance-associated protein 4 is involved in the urinary excretion of hydrochlorothiazide and furosemide. *J Am Soc Nephrol* 18:37-45
 40. Hebert SC, Mount DB, Gamba G (2004) Molecular physiology of cation-coupled Cl-cotransport: the SLC12 family. *Pflugers Arch* 447:580-593
 41. Hoorn EJ, Ellison DH (2012) WNK kinases and the kidney. *Exp Cell Res* 318:1020-1026
 42. Hoorn EJ, Nelson JH, McCormick JA, Ellison DH (2011) The WNK kinase network regulating sodium, potassium, and blood pressure. *J Am Soc Nephrol* 22:605-614
 43. Hoorn EJ, Pisitkun T, Zietse R, Gross P, Frokiaer J, Wang NS, Gonzales PA, Star RA, Knepper MA (2005) Prospects for urinary proteomics: exosomes as a source of urinary biomarkers. *Nephrology (Carlton)* 10:283-290
 44. Hoorn EJ, Walsh SB, McCormick JA, Furstenberg A, Yang CL, Roeschel T, Paliege A, Howie AJ, Conley J, Bachmann S, Unwin RJ, Ellison DH (2011) The calcineurin inhibitor tacrolimus activates the renal sodium chloride cotransporter to cause hypertension. *Nat Med* 17:1304-1309
 45. Hoorn EJ, Walsh SB, McCormick JA, Zietse R, Unwin RJ, Ellison DH (2012) Pathogenesis of calcineurin inhibitor-induced hypertension. *J Nephrol* 25:269-275

46. Hoorn EJ, Walsh SB, Unwin RJ, Ellison DH (2012) Hypertension after kidney transplantation: calcineurin inhibitors increase salt-sensitivity. *J Hypertens* 30:832-833; author reply 833-834
47. Hoorn EJ, Zietse R (2013) Disorders of calcium and magnesium balance: a physiology-based approach. *Pediatr Nephrol* 28:1195-1206
48. Hoorn EJ, Zietse R (2008) Hyponatremia revisited: translating physiology to practice. *Nephron Physiol* 108:p46-59
49. Hoover RS (2011) Angiotensin II: a candidate for an aldosterone-independent mediator of potassium preservation during volume depletion. *Kidney Int* 79:377-379
50. Hoover RS, Poch E, Monroy A, Vazquez N, Nishio T, Gamba G, Hebert SC (2003) N-Glycosylation at two sites critically alters thiazide binding and activity of the rat thiazide-sensitive Na(+):Cl(-) cotransporter. *J Am Soc Nephrol* 14:271-282
51. Hossain Khan MZ, Sohara E, Ohta A, Chiga M, Inoue Y, Isobe K, Wakabayashi M, Oi K, Rai T, Sasaki S, Uchida S (2012) Phosphorylation of Na-Cl cotransporter by OSR1 and SPAK kinases regulates its ubiquitination. *Biochem Biophys Res Commun* 425:456-461
52. Ji W, Foo JN, O'Roak BJ, Zhao H, Larson MG, Simon DB, Newton-Cheh C, State MW, Levy D, Lifton RP (2008) Rare independent mutations in renal salt handling genes contribute to blood pressure variation. *Nat Genet* 40:592-599
53. Kim GH, Masilamani S, Turner R, Mitchell C, Wade JB, Knepper MA (1998) The thiazide-sensitive Na-Cl cotransporter is an aldosterone-induced protein. *Proc Natl Acad Sci U S A* 95:14552-14557
54. Ko B, Joshi LM, Cooke LL, Vazquez N, Musch MW, Hebert SC, Gamba G, Hoover RS (2007) Phorbol ester stimulation of RasGRP1 regulates the sodium-chloride cotransporter by a PKC-independent pathway. *Proc Natl Acad Sci U S A* 104:20120-20125
55. Ko B, Kamsteeg EJ, Cooke LL, Moddes LN, Deen PM, Hoover RS (2010) RasGRP1 stimulation enhances ubiquitination and endocytosis of the sodium-chloride cotransporter. *Am J Physiol Renal Physiol* 299:F300-309
56. Ko B, Mistry AC, Hanson LN, Mallick R, Wynne BM, Thai TL, Bailey JL, Klein JD, Hoover RS (2013) Aldosterone Acutely Stimulates Ncc Activity Via a Spak-Mediated Pathway. *Am J Physiol Renal Physiol*
57. Komers R, Rogers S, Oyama TT, Xu B, Yang CL, McCormick J, Ellison DH (2012) Enhanced phosphorylation of Na(+)-Cl- co-transporter in experimental metabolic syndrome: role of insulin. *Clin Sci (Lond)* 123:635-647
58. Konrad M, Vollmer M, Lemmink HH, van den Heuvel LP, Jeck N, Vargas-Poussou R, Lakings A, Ruf R, Deschenes G, Antignac C, Guay-Woodford L, Knoers NV, Seyberth HW, Feldmann D, Hildebrandt F (2000) Mutations in the chloride channel gene CLCNKB as a cause of classic Bartter syndrome. *J Am Soc Nephrol* 11:1449-1459
59. Kriz W, Bankir L (1988) A standard nomenclature for structures of the kidney. The Renal Commission of the International Union of Physiological Sciences (IUPS). *Kidney Int* 33:1-7
60. Kunau RT, Jr., Weller DR, Webb HL (1975) Clarification of the site of action of chlorothiazide in the rat nephron. *J Clin Invest* 56:401-407
61. Lalioti MD, Zhang J, Volkman HM, Kahle KT, Hoffmann KE, Toka HR, Nelson-Williams C, Ellison DH, Flavell R, Booth CJ, Lu Y, Geller DS, Lifton RP (2006) Wnk4

- controls blood pressure and potassium homeostasis via regulation of mass and activity of the distal convoluted tubule. *Nat Genet* 38:1124-1132
62. Ledeganck KJ, Boulet GA, Horvath CA, Vinckx M, Bogers JJ, Van Den Bossche R, Verpooten GA, De Winter BY (2011) Expression of renal distal tubule transporters TRPM6 and NCC in a rat model of cyclosporine nephrotoxicity and effect of EGF treatment. *Am J Physiol Renal Physiol* 301:F486-493
 63. Leviel F, Hubner CA, Houillier P, Morla L, El Moghrabi S, Brideau G, Hassan H, Parker MD, Kurth I, Kougioumtzes A, Sinning A, Pech V, Riemondy KA, Miller RL, Hummler E, Shull GE, Aronson PS, Doucet A, Wall SM, Chambrey R, Eladari D (2010) The Na⁺-dependent chloride-bicarbonate exchanger SLC4A8 mediates an electroneutral Na⁺ reabsorption process in the renal cortical collecting ducts of mice. *J Clin Invest* 120:1627-1635
 64. Lin SH, Cheng NL, Hsu YJ, Halperin ML (2004) Intrafamilial phenotype variability in patients with Gitelman syndrome having the same mutations in their thiazide-sensitive sodium/chloride cotransporter. *Am J Kidney Dis* 43:304-312
 65. Lin SH, Shiang JC, Huang CC, Yang SS, Hsu YJ, Cheng CJ (2005) Phenotype and genotype analysis in Chinese patients with Gitelman's syndrome. *J Clin Endocrinol Metab* 90:2500-2507
 66. Loffing J, Vallon V, Loffing-Cueni D, Aregger F, Richter K, Pietri L, Bloch-Faure M, Hoenderop JG, Shull GE, Meneton P, Kaissling B (2004) Altered renal distal tubule structure and renal Na⁽⁺⁾ and Ca⁽²⁺⁾ handling in a mouse model for Gitelman's syndrome. *J Am Soc Nephrol* 15:2276-2288
 67. Loffing J, Zecevic M, Feraille E, Kaissling B, Asher C, Rossier BC, Firestone GL, Pearce D, Verrey F (2001) Aldosterone induces rapid apical translocation of ENaC in early portion of renal collecting system: possible role of SGK. *Am J Physiol Renal Physiol* 280:F675-682
 68. Louis-Dit-Picard H, Barc J, Trujillano D, Miserey-Lenkei S, Bouatia-Naji N, Pylypenko O, Beaurain G, Bonnefond A, Sand O, Simian C, Vidal-Petiot E, Soukaseum C, Mandet C, Broux F, Chabre O, Delahousse M, Esnault V, Fiquet B, Houillier P, Bagnis CI, Koenig J, Konrad M, Landais P, Mourani C, Niaudet P, Probst V, Thauvin C, Unwin RJ, Soroka SD, Ehret G, Ossowski S, Caulfield M, International Consortium for Blood P, Bruneval P, Estivill X, Froguel P, Hadchouel J, Schott JJ, Jeunemaitre X (2012) KLHL3 mutations cause familial hyperkalemic hypertension by impairing ion transport in the distal nephron. *Nat Genet* 44:456-460, S451-453
 69. Masilamani S, Wang X, Kim GH, Brooks H, Nielsen J, Nielsen S, Nakamura K, Stokes JB, Knepper MA (2002) Time course of renal Na-K-ATPase, NHE3, NKCC2, NCC, and ENaC abundance changes with dietary NaCl restriction. *Am J Physiol Renal Physiol* 283:F648-657
 70. Mastroianni N, De Fusco M, Zollo M, Arrigo G, Zuffardi O, Bettinelli A, Ballabio A, Casari G (1996) Molecular cloning, expression pattern, and chromosomal localization of the human Na-Cl thiazide-sensitive cotransporter (SLC12A3). *Genomics* 35:486-493
 71. Mayan H, Vered I, Mouallem M, Tzadok-Witkon M, Pazner R, Farfel Z (2002) Pseudohypoaldosteronism type II: marked sensitivity to thiazides, hypercalciuria, normomagnesemia, and low bone mineral density. *J Clin Endocrinol Metab* 87:3248-3254

72. McCormick JA, Ellison DH (2011) The WNKs: atypical protein kinases with pleiotropic actions. *Physiol Rev* 91:177-219
73. McCormick JA, Mutig K, Nelson JH, Saritas T, Hoorn EJ, Yang CL, Rogers S, Curry J, Delpire E, Bachmann S, Ellison DH (2011) A SPAK isoform switch modulates renal salt transport and blood pressure. *Cell Metab* 14:352-364
74. McCormick JA, Nelson JH, Yang CL, Curry JN, Ellison DH (2011) Overexpression of the sodium chloride cotransporter is not sufficient to cause familial hyperkalemic hypertension. *Hypertension* 58:888-894
75. McDonough AA (2010) Mechanisms of proximal tubule sodium transport regulation that link extracellular fluid volume and blood pressure. *Am J Physiol Regul Integr Comp Physiol* 298:R851-861
76. Melnikov S, Mayan H, Uchida S, Holtzman EJ, Farfel Z (2011) Cyclosporine metabolic side effects: association with the WNK4 system. *Eur J Clin Invest* 41:1113-1120
77. Meneton P, Loffing J, Warnock DG (2004) Sodium and potassium handling by the aldosterone-sensitive distal nephron: the pivotal role of the distal and connecting tubule. *Am J Physiol Renal Physiol* 287:F593-601
78. Moreno E, Cristobal PS, Rivera M, Vazquez N, Bobadilla NA, Gamba G (2006) Affinity-defining domains in the Na-Cl cotransporter: a different location for Cl⁻ and thiazide binding. *J Biol Chem* 281:17266-17275
79. Moriguchi T, Urushiyama S, Hisamoto N, Iemura S, Uchida S, Natsume T, Matsumoto K, Shibuya H (2005) WNK1 regulates phosphorylation of cation-chloride-coupled cotransporters via the STE20-related kinases, SPAK and OSR1. *J Biol Chem* 280:42685-42693
80. Morris RG, Hoorn EJ, Knepper MA (2006) Hypokalemia in a mouse model of Gitelman's syndrome. *Am J Physiol Renal Physiol* 290:F1416-1420
81. Moser M, Feig PU (2009) Fifty years of thiazide diuretic therapy for hypertension. *Arch Intern Med* 169:1851-1856
82. Mu S, Shimosawa T, Ogura S, Wang H, Uetake Y, Kawakami-Mori F, Marumo T, Yatomi Y, Geller DS, Tanaka H, Fujita T (2011) Epigenetic modulation of the renal beta-adrenergic-WNK4 pathway in salt-sensitive hypertension. *Nat Med* 17:573-580
83. Mutig K, Saritas T, Uchida S, Kahl T, Borowski T, Paliege A, Bohlick A, Bleich M, Shan Q, Bachmann S (2010) Short-term stimulation of the thiazide-sensitive Na⁺-Cl⁻ cotransporter by vasopressin involves phosphorylation and membrane translocation. *Am J Physiol Renal Physiol* 298:F502-509
84. Nguyen MT, Lee DH, Delpire E, McDonough AA (2013) Differential Regulation of Na⁺ Transporters along Nephron during AngII-Dependent Hypertension: Distal Stimulation Counteracted by Proximal Inhibition. *Am J Physiol Renal Physiol*
85. Nijenhuis T, Vallon V, van der Kemp AW, Loffing J, Hoenderop JG, Bindels RJ (2005) Enhanced passive Ca²⁺ reabsorption and reduced Mg²⁺ channel abundance explains thiazide-induced hypocalciuria and hypomagnesemia. *J Clin Invest* 115:1651-1658
86. Novello FC, Sprague JM (1957) Benzothiadiazine dioxides as novel diuretics. *J Am Chem Soc* 79:2028- 2029
87. O'Reilly M, Marshall E, Macgillivray T, Mittal M, Xue W, Kenyon CJ, Brown RW (2006) Dietary electrolyte-driven responses in the renal WNK kinase pathway in vivo. *J Am Soc Nephrol* 17:2402-2413

88. Ohta A, Rai T, Yui N, Chiga M, Yang SS, Lin SH, Sohara E, Sasaki S, Uchida S (2009) Targeted disruption of the Wnk4 gene decreases phosphorylation of Na-Cl cotransporter, increases Na excretion and lowers blood pressure. *Hum Mol Genet* 18:3978-3986
89. Ohta A, Schumacher FR, Mehellou Y, Johnson C, Knebel A, Macartney TJ, Wood NT, Alessi DR, Kurz T (2013) The CUL3-KLHL3 E3 ligase complex mutated in Gordon's hypertension syndrome interacts with and ubiquitylates WNK isoforms: disease-causing mutations in KLHL3 and WNK4 disrupt interaction. *Biochem J* 451:111-122
90. Pacheco-Alvarez D, Vazquez N, Castaneda-Bueno M, de-Los-Heros P, Cortes-Gonzalez C, Moreno E, Meade P, Bobadilla NA, Gamba G (2012) WNK3-SPAK interaction is required for the modulation of NCC and other members of the SLC12 family. *Cell Physiol Biochem* 29:291-302
91. Pedersen NB, Hofmeister MV, Rosenbaek LL, Nielsen J, Fenton RA (2010) Vasopressin induces phosphorylation of the thiazide-sensitive sodium chloride cotransporter in the distal convoluted tubule. *Kidney Int* 78:160-169
92. Pickkers P, Hughes AD, Russel FG, Thien T, Smits P (1998) Thiazide-induced vasodilation in humans is mediated by potassium channel activation. *Hypertension* 32:1071-1076
93. Pisitkun T, Shen RF, Knepper MA (2004) Identification and proteomic profiling of exosomes in human urine. *Proc Natl Acad Sci U S A* 101:13368-13373
94. Rafiqi FH, Zuber AM, Glover M, Richardson C, Fleming S, Jovanovic S, Jovanovic A, O'Shaughnessy KM, Alessi DR (2010) Role of the WNK-activated SPAK kinase in regulating blood pressure. *EMBO Mol Med* 2:63-75
95. Renfro JL (1977) Interdependence of Active Na⁺ and Cl⁻ transport by the isolated urinary bladder of the teleost, *Pseudopleuronectes americanus*. *J Exp Zool* 199:383-390
96. Renfro JL (1975) Water and ion transport by the urinary bladder of the teleost *Pseudopleuronectes americanus*. *Am J Physiol* 228:52-61
97. Rinehart J, Kahle KT, de Los Heros P, Vazquez N, Meade P, Wilson FH, Hebert SC, Gimenez I, Gamba G, Lifton RP (2005) WNK3 kinase is a positive regulator of NKCC2 and NCC, renal cation-Cl⁻ cotransporters required for normal blood pressure homeostasis. *Proc Natl Acad Sci U S A* 102:16777-16782
98. Ring AM, Leng Q, Rinehart J, Wilson FH, Kahle KT, Hebert SC, Lifton RP (2007) An SGK1 site in WNK4 regulates Na⁺ channel and K⁺ channel activity and has implications for aldosterone signaling and K⁺ homeostasis. *Proc Natl Acad Sci U S A* 104:4025-4029
99. Ronzaud C, Loffing-Cueni D, Hausel P, Debonneville A, Malsure SR, Fowler-Jaeger N, Boase NA, Perrier R, Maillard M, Yang B, Stokes JB, Koesters R, Kumar S, Hummler E, Loffing J, Staub O (2013) Renal tubular NEDD4-2 deficiency causes NCC-mediated salt-dependent hypertension. *J Clin Invest* 123:657-665
100. Rozansky DJ, Cornwall T, Subramanya AR, Rogers S, Yang YF, David LL, Zhu X, Yang CL, Ellison DH (2009) Aldosterone mediates activation of the thiazide-sensitive Na-Cl cotransporter through an SGK1 and WNK4 signaling pathway. *J Clin Invest* 119:2601-2612
101. San-Cristobal P, Pacheco-Alvarez D, Richardson C, Ring AM, Vazquez N, Rafiqi FH, Chari D, Kahle KT, Leng Q, Bobadilla NA, Hebert SC, Alessi DR, Lifton RP, Gamba G (2009) Angiotensin II signaling increases activity of the renal Na-Cl cotransporter through a WNK4-SPAK-dependent pathway. *Proc Natl Acad Sci U S A* 106:4384-4389

102. San-Cristobal P, Ponce-Coria J, Vazquez N, Bobadilla NA, Gamba G (2008) WNK3 and WNK4 amino-terminal domain defines their effect on the renal Na⁺-Cl⁻ cotransporter. *Am J Physiol Renal Physiol* 295:F1199-1206
103. Sandberg MB, Maunsbach AB, McDonough AA (2006) Redistribution of distal tubule Na⁺-Cl⁻ cotransporter (NCC) in response to a high-salt diet. *Am J Physiol Renal Physiol* 291:F503-508
104. Saritas T, Borschewski A, McCormick JA, Paliege A, Dathe C, Uchida S, Terker A, Himmerkus N, Bleich M, Demaretz S, Laghmani K, Delpire E, Ellison DH, Bachmann S, Mutig K (2013) SPAK differentially mediates vasopressin effects on sodium cotransporters. *J Am Soc Nephrol* 24:407-418
105. Schultheis PJ, Lorenz JN, Meneton P, Nieman ML, Riddle TM, Flagella M, Duffy JJ, Doetschman T, Miller ML, Shull GE (1998) Phenotype resembling Gitelman's syndrome in mice lacking the apical Na⁺-Cl⁻ cotransporter of the distal convoluted tubule. *J Biol Chem* 273:29150-29155
106. Shibata S, Zhang J, Puthumana J, Stone KL, Lifton RP (2013) Kelch-like 3 and Cullin 3 regulate electrolyte homeostasis via ubiquitination and degradation of WNK4. *Proc Natl Acad Sci U S A* 110:7838-7843
107. Simon DB, Nelson-Williams C, Bia MJ, Ellison D, Karet FE, Molina AM, Vaara I, Iwata F, Cushner HM, Koolen M, Gainza FJ, Gitleman HJ, Lifton RP (1996) Gitelman's variant of Bartter's syndrome, inherited hypokalaemic alkalosis, is caused by mutations in the thiazide-sensitive Na-Cl cotransporter. *Nat Genet* 12:24-30
108. Sohara E, Rai T, Yang SS, Ohta A, Naito S, Chiga M, Nomura N, Lin SH, Vandewalle A, Ohta E, Sasaki S, Uchida S (2011) Acute insulin stimulation induces phosphorylation of the Na-Cl cotransporter in cultured distal mpkDCT cells and mouse kidney. *PLoS One* 6:e24277
109. Song J, Hu X, Riazi S, Tiwari S, Wade JB, Ecelbarger CA (2006) Regulation of blood pressure, the epithelial sodium channel (ENaC), and other key renal sodium transporters by chronic insulin infusion in rats. *Am J Physiol Renal Physiol* 290:F1055-1064
110. Sorensen MV, Grossmann S, Roesinger M, Gresko N, Todkar AP, Barmettler G, Ziegler U, Odermatt A, Loffing-Cueni D, Loffing J (2013) Rapid dephosphorylation of the renal sodium chloride cotransporter in response to oral potassium intake in mice. *Kidney Int* 83:811-824
111. Stokes JB (1984) Sodium chloride absorption by the urinary bladder of the winter flounder. A thiazide-sensitive, electrically neutral transport system. *J Clin Invest* 74:7-16
112. Subramanya AR, Yang CL, McCormick JA, Ellison DH (2006) WNK kinases regulate sodium chloride and potassium transport by the aldosterone-sensitive distal nephron. *Kidney Int* 70:630-634
113. Tseng MH, Yang SS, Hsu YJ, Fang YW, Wu CJ, Tsai JD, Hwang DY, Lin SH (2012) Genotype, phenotype, and follow-up in Taiwanese patients with salt-losing tubulopathy associated with SLC12A3 mutation. *J Clin Endocrinol Metab* 97:E1478-1482
114. Vallon V, Rieg T, Ahn SY, Wu W, Eraly SA, Nigam SK (2008) Overlapping in vitro and in vivo specificities of the organic anion transporters OAT1 and OAT3 for loop and thiazide diuretics. *Am J Physiol Renal Physiol* 294:F867-873
115. Vallon V, Schroth J, Lang F, Kuhl D, Uchida S (2009) Expression and phosphorylation of the Na⁺-Cl⁻ cotransporter NCC in vivo is regulated by dietary salt, potassium, and SGK1. *Am J Physiol Renal Physiol* 297:F704-712

116. van Angelen AA, Glaudemans B, van der Kemp AW, Hoenderop JG, Bindels RJ (2013) Cisplatin-induced injury of the renal distal convoluted tubule is associated with hypomagnesaemia in mice. *Nephrol Dial Transplant* 28:879-889
117. van der Lubbe N, Jansen PM, Salih M, Fenton RA, van den Meiracker AH, Danser AH, Zietse R, Hoorn EJ (2012) The phosphorylated sodium chloride cotransporter in urinary exosomes is superior to prostaticin as a marker for aldosteronism. *Hypertension* 60:741-748
118. van der Lubbe N, Lim CH, Fenton RA, Meima ME, Jan Danser AH, Zietse R, Hoorn EJ (2011) Angiotensin II induces phosphorylation of the thiazide-sensitive sodium chloride cotransporter independent of aldosterone. *Kidney Int* 79:66-76
119. van der Lubbe N, Lim CH, Meima ME, van Veghel R, Rosenbaek LL, Mutig K, Danser AH, Fenton RA, Zietse R, Hoorn EJ (2012) Aldosterone does not require angiotensin II to activate NCC through a WNK4-SPAK-dependent pathway. *Pflugers Arch* 463:853-863
120. Van der Lubbe N, Moes A, Rosenbaek LL, Schoep S, Meima ME, Danser AH, Fenton RA, Zietse R, Hoorn EJ (2013) Potassium-induced natriuresis is preserved during sodium depletion and accompanied by inhibition of the sodium chloride cotransporter. In revision
121. van der Lubbe N, Moes AD, Rosenbaek LL, Schoep S, Meima ME, Danser AH, Fenton RA, Zietse R, Hoorn EJ (2013) K⁺-induced natriuresis is preserved during Na⁺ depletion and accompanied by inhibition of the Na⁺-Cl⁻ cotransporter. *Am J Physiol Renal Physiol* 305:F1177-1188
122. van der Lubbe N, Zietse R, Hoorn EJ (2013) Effects of angiotensin II on kinase-mediated sodium and potassium transport in the distal nephron. *Curr Opin Nephrol Hypertens* 22:120-126
123. Vargas-Poussou R, Dahan K, Kahila D, Venisse A, Riveira-Munoz E, Debaix H, Grisart B, Bridoux F, Unwin R, Moulin B, Haymann JP, Vantyghem MC, Rigotherier C, Dussol B, Godin M, Nivet H, Dubourg L, Tack I, Gimenez-Roqueplo AP, Houillier P, Blanchard A, Devuyst O, Jeunemaitre X (2011) Spectrum of mutations in Gitelman syndrome. *J Am Soc Nephrol* 22:693-703
124. Velazquez H, Bartiss A, Bernstein P, Ellison DH (1996) Adrenal steroids stimulate thiazide-sensitive NaCl transport by rat renal distal tubules. *Am J Physiol* 270:F211-219
125. Verlander JW, Tran TM, Zhang L, Kaplan MR, Hebert SC (1998) Estradiol enhances thiazide-sensitive NaCl cotransporter density in the apical plasma membrane of the distal convoluted tubule in ovariectomized rats. *J Clin Invest* 101:1661-1669
126. Wagner CA, Loffing-Cueni D, Yan Q, Schulz N, Fakitsas P, Carrel M, Wang T, Verrey F, Geibel JP, Giebisch G, Hebert SC, Loffing J (2008) Mouse model of type II Bartter's syndrome. II. Altered expression of renal sodium- and water-transporting proteins. *Am J Physiol Renal Physiol* 294:F1373-1380
127. Wagner CA, Ott M, Klingel K, Beck S, Melzig J, Friedrich B, Wild KN, Broer S, Moschen I, Albers A, Waldegger S, Tummler B, Egan ME, Geibel JP, Kandolf R, Lang F (2001) Effects of the serine/threonine kinase SGK1 on the epithelial Na(+) channel (ENaC) and CFTR: implications for cystic fibrosis. *Cell Physiol Biochem* 11:209-218
128. Wakabayashi M, Mori T, Isobe K, Sohara E, Susa K, Araki Y, Chiga M, Kikuchi E, Nomura N, Mori Y, Matsuo H, Murata T, Nomura S, Asano T, Kawaguchi H, Nonoyama S, Rai T, Sasaki S, Uchida S (2013) Impaired KLHL3-mediated ubiquitination of WNK4 causes human hypertension. *Cell Rep* 3:858-868

129. Wei Y, Zavilowitz B, Satlin LM, Wang WH (2007) Angiotensin II inhibits the ROMK-like small conductance K channel in renal cortical collecting duct during dietary potassium restriction. *J Biol Chem* 282:6455-6462
130. Welling PA, Chang YP, Delpire E, Wade JB (2010) Multigene kinase network, kidney transport, and salt in essential hypertension. *Kidney Int* 77:1063-1069
131. Wilson FH, Disse-Nicodeme S, Choate KA, Ishikawa K, Nelson-Williams C, Desitter I, Gunel M, Milford DV, Lipkin GW, Achard JM, Feely MP, Dussol B, Berland Y, Unwin RJ, Mayan H, Simon DB, Farfel Z, Jeunemaitre X, Lifton RP (2001) Human hypertension caused by mutations in WNK kinases. *Science* 293:1107-1112
132. Wu G, Peng JB (2013) Disease-causing mutations in KLHL3 impair its effect on WNK4 degradation. *FEBS Lett* 587:1717-1722
133. Xu JZ, Hall AE, Peterson LN, Bienkowski MJ, Eessalu TE, Hebert SC (1997) Localization of the ROMK protein on apical membranes of rat kidney nephron segments. *Am J Physiol* 273:F739-748
134. Yang CL, Angell J, Mitchell R, Ellison DH (2003) WNK kinases regulate thiazide-sensitive Na-Cl cotransport. *J Clin Invest* 111:1039-1045
135. Yang CL, Zhu X, Ellison DH (2007) The thiazide-sensitive Na-Cl cotransporter is regulated by a WNK kinase signaling complex. *J Clin Invest* 117:3403-3411
136. Yang CL, Zhu X, Wang Z, Subramanya AR, Ellison DH (2005) Mechanisms of WNK1 and WNK4 interaction in the regulation of thiazide-sensitive NaCl cotransport. *J Clin Invest* 115:1379-1387
137. Yang LE, Sandberg MB, Can AD, Pihakaski-Maunsbach K, McDonough AA (2008) Effects of dietary salt on renal Na⁺ transporter subcellular distribution, abundance, and phosphorylation status. *Am J Physiol Renal Physiol* 295:F1003-1016
138. Yang SS, Fang YW, Tseng MH, Chu PY, Yu IS, Wu HC, Lin SW, Chau T, Uchida S, Sasaki S, Lin YF, Sytwu HK, Lin SH (2013) Phosphorylation regulates NCC stability and transporter activity in vivo. *J Am Soc Nephrol* 24:1587-1597
139. Yang SS, Lo YF, Wu CC, Lin SW, Yeh CJ, Chu P, Sytwu HK, Uchida S, Sasaki S, Lin SH (2010) SPAK-knockout mice manifest Gitelman syndrome and impaired vasoconstriction. *J Am Soc Nephrol* 21:1868-1877
140. Yue P, Sun P, Lin DH, Pan C, Xing W, Wang W (2011) Angiotensin II diminishes the effect of SGK1 on the WNK4-mediated inhibition of ROMK1 channels. *Kidney Int* 79:423-431
141. Zhou B, Wang D, Feng X, Zhang Y, Wang Y, Zhuang J, Zhang X, Chen G, Delpire E, Gu D, Cai H (2012) WNK4 inhibits NCC protein expression through MAPK ERK1/2 signaling pathway. *Am J Physiol Renal Physiol* 302:F533-539
142. Zhou B, Zhuang J, Gu D, Wang H, Cebotaru L, Guggino WB, Cai H (2010) WNK4 enhances the degradation of NCC through a sortilin-mediated lysosomal pathway. *J Am Soc Nephrol* 21:82-92

Figure 1
[Click here to download high resolution image](#)




Figure
[Click here to download high resolution image](#)




Figure 3

[Click here to download high resolution image](#)

